

# Reciprocal Regulation of Brain and Muscle Arnt-Like Protein 1 and Peroxisome Proliferator-Activated Receptor $\alpha$ Defines a Novel Positive Feedback Loop in the Rodent Liver Circadian Clock

Laurence Canaple, Juliette Rambaud, Ouria Dkhissi-Benyahya, Béatrice Rayet, Nguan Soon Tan, Liliane Michalik, Franck Delaunay, Walter Wahli, and Vincent Laudet

*Structure and Evolution of Nuclear Receptors (L.C., J.R., V.L.), Centre National de la Recherche Scientifique (CNRS) Unité Mixte de Recherche 5161, Institut Fédératif de Recherche (IFR) 128 BioSciences Lyon-Gerland, Laboratoire de Biologie Moléculaire de la Cellule, Ecole Normale Supérieure, 69364 Lyon cedex 07, France; Institut National de la Santé et de la Recherche Médicale Unité 371 (O.D.-B.), Laboratoire Cerveau et Vision, IFR 19-Université Claude Bernard Lyon 1, 69675 Bron, France; Université de Nice Sophia Antipolis CNRS Formation de Recherche en Evolution (B.R., F.D.), 06108 Nice cedex 2, France; and Center for Integrative Genomics (N.S.T., L.M., W.W.), University of Lausanne, CH-1015 Lausanne, Switzerland*

Recent evidence has emerged that peroxisome proliferator-activated receptor  $\alpha$  (PPAR $\alpha$ ), which is largely involved in lipid metabolism, can play an important role in connecting circadian biology and metabolism. In the present study, we investigated the mechanisms by which PPAR $\alpha$  influences the pacemakers acting in the central clock located in the suprachiasmatic nucleus and in the peripheral oscillator of the liver. We demonstrate that PPAR $\alpha$  plays a specific role in the peripheral circadian control because it is required to maintain the circadian rhythm of the master clock gene brain and

muscle Arnt-like protein 1 (*bmal1*) *in vivo*. This regulation occurs via a direct binding of PPAR $\alpha$  on a potential PPAR $\alpha$  response element located in the *bmal1* promoter. Reversely, BMAL1 is an upstream regulator of PPAR $\alpha$  gene expression. We further demonstrate that fenofibrate induces circadian rhythm of clock gene expression in cell culture and up-regulates hepatic *bmal1* *in vivo*. Together, these results provide evidence for an additional regulatory feedback loop involving BMAL1 and PPAR $\alpha$  in peripheral clocks. (*Molecular Endocrinology* 20: 1715–1727, 2006)

CIRCADIAN RHYTHMS ENABLE numerous organisms to adapt to daily environmental changes such as light, temperature, and social communication and serve to synchronize multiple molecular, biochemical, physiological, and behavioral processes. Circadian rhythms persist with an approximate 24-h periodicity even in temporally isolated subjects, indicating the presence of an autonomous time-keeping system called circadian clock. In mammals, circadian rhythms are generated by the main pacemaker located in the suprachiasmatic nucleus (SCN) of the hypothalamus (1). To ensure that internal time coincides with environmental time, the clock must be adjusted, a process known as entrainment. In mammals, light re-

ceived by the eyes synchronizes the oscillator through the retino-hypothalamic tract and hence synchronizes the behavior of the organism with the daily 24-h light-dark (LD) cycle (for review, see Refs. 2–6).

In addition to the SCN, other peripheral tissues such as liver, heart, kidney (7, 8), as well as isolated cells (9) express clock genes giving rise to circadian rhythms with a different phase from that observed in the SCN. Interestingly, these peripheral clocks can be reset by alternative routes independently of the SCN, for example by forced change of feeding time (10, 11). Several lines of evidence suggest that the peripheral circadian clocks are not SCN independent but require inputs from the SCN to drive the rhythmicity and ensure an ordered response of the organism to environmental changes (12, 13). Thus, the SCN is believed to coordinate rhythms in the brain and body via a combination of neural and humoral diffusible and synaptic signals (7, 14, 15).

Genetic analyses have identified master clock genes such as *clock*, *bmal1*, *period* (*per1*, 2), and *cryptochrome* genes (*cry1*, 2), as well as the orphan nuclear receptor genes, *rev-erb $\alpha$*  and *ror $\alpha$*  (16, 17). Other transcription factors functioning in the circadian regulation of gene expression, including DBP-related factors

## First Published Online March 23, 2006

Abbreviations: BMAL1, Brain and muscle Arnt-like protein 1; ChIP, chromatin immunoprecipitation; CT, circadian time; DD, constant darkness; LD, light-dark; 12L:12D, 12-h light, 12-h dark; PPAR $\alpha$ , peroxisome proliferator-activated receptor  $\alpha$ ; PPRE, peroxisome proliferator response element; PSG, penicillin/streptomycin/glutamine; SCN, suprachiasmatic nucleus; WT, wild type; ZT, Zeitgeber time.

**Molecular Endocrinology** is published monthly by The Endocrine Society (<http://www.endo-society.org>), the foremost professional society serving the endocrine community.

(18), Rev-erb $\beta$ , ROR $\beta$  and  $\gamma$  (19) have also been identified. The clock mechanism mainly involves an integrated network of interacting self-sustained transcriptional-translational feedback loops, composed of positive and negative regulators, which drive their own rhythmic expression and the one of clock-controlled genes to perform a fine tuning of circadian gene expression (20).

Recent reports have highlighted the interplay between circadian oscillators, metabolism, and physiology. Whereas genes involved in the glucose and lipid metabolism are known to exhibit circadian variations (21–23), molecular studies have revealed a critical role for *bmal1* and *clock* genes in regulating glucose homeostasis (24) and lipid metabolism (25–28). Moreover, the cross talk between ROR $\alpha$  and Rev-erb $\alpha$  was shown to be physiologically important for the control of cholesterol and triglyceride metabolism (29–31). In turn, the peripheral clocks can be coordinately regulated by multiple circulating factors, which are affected by the metabolic status of the organism. Indeed, glucose, one of the major food metabolites that exhibits a plasma diurnal rhythm, is a direct resetting signal in cultured cells by down-regulating *per1* and *per2* RNA levels (32). The levels of glucose-regulated hormones such as insulin or glucagon immediately up-regulate *per1* and *per2* expression (13). In addition, other studies have revealed an important role of glucocorticoids and retinoids in the resetting of peripheral clocks (33, 34). Although the elucidation of the mechanisms that govern the connection between metabolism and circadian clock has just begun, it appears that several members of the nuclear receptor family are involved in this pathway.

Evidence has emerged that peroxisome proliferator-activated receptor  $\alpha$  (PPAR $\alpha$ ), a member of the nuclear receptor superfamily that regulates the expression of numerous genes involved in lipid metabolism and energy homeostasis, can play a role in the normal circadian regulation. First, PPAR $\alpha$  has been identified as a circadian clock-controlled gene with a diurnal rhythm at the mRNA and protein levels in rats and mice in many peripheral organs such as liver, heart, kidney, and to a lesser extent in the SCN, where the central pacemaker is located (7, 35, 36). This circadian expression of PPAR $\alpha$  may be in part controlled by hormonal factors, because insulin and glucocorticoids regulate its mRNA expression (35, 37–39). A recent study has also shown that the circadian expression of PPAR $\alpha$  mRNA is regulated by the peripheral oscillators in a CLOCK-dependent manner (27). Second, because daily variations in lipogenic and cholesterogenic gene expression are attenuated or abolished in mice in which the PPAR $\alpha$  gene has been disrupted, PPAR $\alpha$  may be an important mediator for the circadian regulation of lipid metabolism (40, 41). It is now believed that PPAR $\alpha$  has a wider general role in transducing hormone messages involved in dietary status (42). These observations thus suggest that PPAR $\alpha$  may be required in the control of circadian food-dependent

fluctuations in gene expression. Third, PPAR $\alpha$  is connected to the regulation of other nuclear hormone receptors such as Rev-erb $\alpha$ , because fenofibrate, a PPAR $\alpha$  agonist, induces human and rat *rev-erb $\alpha$*  expression in liver through the direct binding of PPAR $\alpha$  on an atypical DR2 element located in the *rev-erb $\alpha$*  promoter (30, 43). It was recently shown that CLOCK plays an important role in lipid homeostasis by regulating the circadian transactivation of potential PPAR $\alpha$  response element (PPRE)-controlled target genes (26) and of PPAR $\alpha$  gene itself via an E-box rich region *in vivo* and *in vitro* (27). Fourth, the partner of PPAR $\alpha$ , RXR $\alpha$ , interacts with CLOCK protein in a ligand-dependent manner and inhibits CLOCK/brain and muscle Arnt-like protein 1 (BMAL1)-dependent activation via an E-box element (34). Other results have also suggested that PPAR $\alpha$  deficiency disturbs the normal circadian regulation of certain SREBP-sensitive genes in the liver (40, 44).

Despite much evidence supporting a role of PPAR $\alpha$  in metabolic control and energy homeostasis (45) and the accumulation of data connecting metabolism and circadian biology (3, 46, 47), little is known concerning the influence of PPAR $\alpha$  on the circadian clock. In the present study, we investigated the mechanisms by which PPAR $\alpha$  can influence the pacemakers acting in the SCN and in the liver. We report that PPAR $\alpha$ -deficient (PPAR $\alpha$ <sup>-/-</sup>) mice present similar locomotor activity with wild-type (WT) mice without any molecular alteration of clock gene expression in the SCN. Interestingly, we show for the first time that PPAR $\alpha$  is a direct regulator of *bmal1* expression in liver via its direct binding on a PPRE located on the *bmal1* promoter. This regulation is required to maintain the normal circadian oscillation of *bmal1* *in vivo*. Stressing the importance of the regulatory pathway that exists between PPAR $\alpha$  and the peripheral clock genes, we show that fenofibrate up-regulates *bmal1* gene expression in murine liver and induces circadian rhythm of clock gene expression in cell culture in a PPAR $\alpha$ -dependent manner. Reversely, we also observe that BMAL1 is an upstream regulator of the PPAR $\alpha$  gene expression. Taken together, our data implicate PPAR $\alpha$  in a new regulatory loop that controls peripheral circadian clocks.


## RESULTS

### PPAR $\alpha$ <sup>-/-</sup> Mice Display Normal Circadian Locomotor Activity and Clock Gene Expression in the SCN

To evaluate the influence of PPAR $\alpha$  on the function of the central circadian oscillator, behavioral analysis of the circadian rhythm was carried out using PPAR $\alpha$ <sup>-/-</sup> and WT littermate mice. Animals were first synchronized for 2 wk to a 12-h light, 12-h dark cycle (12L:12D). Under these light conditions, WT and PPAR $\alpha$ <sup>-/-</sup> mice entrained normally and consolidated their loco-

motor activity to the dark period of the LD cycle (Fig. 1A). No difference in the total amount of daily activity was observed between WT and  $PPAR\alpha^{-/-}$  mice (Fig. 1B). When placed in constant darkness, the  $PPAR\alpha$ -deficient mice do not display an arrhythmic behavior with endogenous period similar between both genotypes ( $24.11 \pm 0.11$  and  $24.12 \pm 0.08$  h, respectively, in  $PPAR\alpha$ -deficient and WT mice).

To determine whether the inactivation of  $PPAR\alpha$  can alter the master oscillator in the SCN at the molecular level, we further compared the circadian expression profiles of clock genes in the SCN of WT and  $PPAR\alpha$  mutant mice (Fig. 1C). There is no significant difference in the amplitude and in the phase of the circadian expression of the tested clock genes (*bmal1*, *per2*, *per3*, *cry2*, and *rev-erb $\alpha$* ). This clearly suggests that



**Fig. 1.** Entrainment and Free Running Locomotor Activity of WT and  $PPAR\alpha^{-/-}$  Mice

A, Representative actograms of locomotor activity for WT and  $PPAR\alpha^{-/-}$  mice under 12L:12D cycle and DD conditions. After 2 wk of entrainment under a 12L:12D cycle, mice were placed in DD. Horizontal bar at the top of each actogram depicts the lighting conditions of LD cycles. Time spans in darkness are marked by gray shadowing. B, Twenty-four-hour profiles of spontaneous locomotor activity of WT ( $\square$ ) and  $PPAR\alpha$ -deficient mice ( $\blacksquare$ ). The distribution of activity was determined every 12 h during LD cycle. Shadowed areas indicate the dark period. Results are expressed as the means  $\pm$  SEM of values from eight animals per group. C, Circadian expression of *bmal1*, *per2*, *per3*, *cry2*, and *rev-erb $\alpha$*  mRNAs in SCN of WT ( $\blacklozenge$ ) and  $PPAR\alpha^{-/-}$  mice ( $\blacksquare$ ). Real-time PCR was used to determine transcript levels at four circadian times (CT1, CT8, CT13, CT21). Transcript levels are displayed as relative quantity (RQ) after normalization to the noncyclic 36B4 expression levels in the same sample. Results are expressed as the means  $\pm$  SEM of values of two independent experiments, each realized with three animals for both genotypes at each time point. Statistically significant differences between WT and  $PPAR\alpha$ -deficient mice are indicated by an asterisk ( $P < 0.05$ ).

there is no major molecular effect of the PPAR $\alpha$  deletion at the central clock level and that the expression of PPAR $\alpha$  in the SCN is not essential for the basal maintenance of the central circadian timing system.


### PPAR $\alpha$ Is Required to Maintain the Amplitude of the Circadian Expression of *bmal1* in the Murine Liver

To evaluate the role of PPAR $\alpha$  on the circadian system of the liver (a peripheral clock), where it is mainly expressed, we analyzed the circadian expression of several clock genes in liver isolated from WT and PPAR $\alpha$  mutant mice. As already reported, PPAR $\alpha$  expression effectively follows a circadian rhythm in peripheral tissues such as liver, kidney, and muscle (data not shown). Figure 2 shows that all the clock genes tested (*bmal1*, *per1*, *per2*, *per3*, *cry2*, and *rev-erb $\alpha$* ) are expressed in a circadian manner with no modification in the phase of their rhythm between both genotypes. By contrast, the amplitudes of *bmal1* and *per3* expression are drastically affected in PPAR $\alpha$ -deficient mice by comparison with the WT, with a significant decrease at circadian time 1 (CT1) and CT21 for *bmal1* (where CT0 is subjective day beginning at 0700 h, and CT12 is subjective night beginning at 1900 h) and an increase at CT8 for *per3*. These data suggest that PPAR $\alpha$  does not influence the phase synchronization

properties of the liver clock but affects the amplitude of two major clock genes *bmal1* and *per3*.


### Food-Induced Phase Resetting Entrains Circadian PPAR $\alpha$ Expression and Is Globally Maintained in Liver of PPAR $\alpha$ <sup>-/-</sup> Mice

Although the day-night cycle is the most obvious time cue, animals can also respond to other synchronizing signals, and feeding time appears to be a potent temporal cue, or Zeitgeber, for the liver clock (10, 11). According to PPAR $\alpha$  role in the hepatic lipid metabolism during starvation, we hypothesized its potential role in food phase resetting of the liver clock. To examine whether the daily feeding time can affect the phase of the PPAR $\alpha$  circadian expression in the liver, WT mice were fed for 2 wk exclusively during the day or during the night. As expected, mice fed exclusively during the day displayed an inversed phase in circadian hepatic expression of *bmal1*, *per1*, *per3*, and *rev-erb $\alpha$*  gene by comparison to mice fed only during the night. Similarly, feeding during the day entirely inversed the phase of liver PPAR $\alpha$  expression (Fig. 3A), a result that is in agreement with the demonstration that PPAR $\alpha$  is a clock-controlled gene in the liver (7, 36). Control mice fed only during the night displayed a similar phase of hepatic clock gene expression and PPAR $\alpha$  than mice fed *ad libitum*.



**Fig. 2.** Circadian Expression of *bmal1*, *per1*, *per2*, *per3*, *cry2*, and *rev-erb $\alpha$*  mRNAs in Liver of WT (◆) and PPAR $\alpha$ <sup>-/-</sup> Mice (■) using Real-Time PCR

Transcript level values are expressed as relative quantity (RQ) after normalization to the corresponding noncyclic 36B4 expression levels. Results are shown as the mean  $\pm$  SEM of values of two independent experiments, each with three animals for both genotypes at each time point. There were significant variations between the two genotypes at CT1 and CT21 for *bmal1* and CT8 for *per3* as indicated by an asterisk ( $P < 0.01$ ).



**Fig. 3.** Circadian Gene Expression in Liver after Restricted Feeding in WT and PPAR $\alpha$ <sup>-/-</sup> Mice

A, PPAR $\alpha$  expression in the liver of food-entrained WT mice (daytime (-♦-) or nighttime feeding (■)). B, Circadian accumulation of *bmal1*, *per1*, *per3*, and *rev-erb $\alpha$*  mRNAs in liver of WT and PPAR $\alpha$ <sup>-/-</sup> mice as a function of daytime (♦) or nighttime feeding (■). Transcript levels are displayed as relative quantity (RQ) after normalization to the corresponding noncyclic 36B4 expression levels. The presented values are expressed as means  $\pm$  SEM of duplicates of the same reaction for six different mice per genotype.

Next, we studied whether feeding time can also reset the phase of *bmal1*, *per1*, *per3*, and *rev-erb $\alpha$*  expression in the absence of PPAR $\alpha$  (Fig. 3B). Except the *bmal1* expression, which was refractory to reset-

ting, *per1*, *per3*, and *rev-erb $\alpha$*  genes showed an inverted rhythm of their expression in the liver after daytime-restricted feeding compared with nighttime feeding in the PPAR $\alpha$  knockout context. The present

results show that feeding time can reset the expression of *per1*, *per3*, and *rev-erb $\alpha$*  in the liver of *PPAR $\alpha$ <sup>-/-</sup>* mice and suggest that *bmal1* expression might be controlled by PPAR $\alpha$ .

### Fenofibrate Induces Expression of Clock Genes in Rat-1 Fibroblasts and Up-Regulates *bmal1* Gene Expression in Liver

Because PPAR $\alpha$  presents a circadian expression in liver, we first studied whether a shock with a serum-rich medium is able to induce an oscillation of PPAR $\alpha$  in the well-established *in vitro* model Rat-1 fibroblasts (9, 13, 48–50). As shown in Fig. 4A, PPAR $\alpha$  gene expression is induced by a serum shock in Rat-1 fibroblasts with a maximal level reached 12 h after the beginning of the treatment in comparison with the control. The induced-oscillating expressions of *rev-erb $\alpha$*  and *cry1* after serum shock were in accordance with previously reported data (9, 51). In addition, *rev-erb $\alpha$*  expression is delayed (peak at 16 h after the serum shock treatment) compared with PPAR $\alpha$  expression, which suggests that in Rat-1 fibroblasts PPAR $\alpha$  induces *rev-erb $\alpha$*  expression. Similarly to these observations, it seems likely that serum largely participates in the synchronization of the circadian oscillation of PPAR $\alpha$  mRNA in fibroblast cultures.

Fibrates are well-known activators of PPAR $\alpha$  expression and are classical drugs used in the treatment of dyslipidemias. Using the same *in vitro* model, we then evaluated whether fenofibrate can induce expression of PPAR $\alpha$  gene. The addition of fenofibrate into the serum-free medium triggered a rhythmic expression of PPAR $\alpha$  peaking 12–14 h as observed after a serum treatment (Fig. 4B). Fenofibrate induced a PPAR $\alpha$  expression 6-fold higher than serum (Fig. 4, A and B).

Fenofibrate can also induce a rhythmic expression of clock genes. The temporal induction by fenofibrate of *cry1*, *rev-erb $\alpha$* , and *bmal1* expression into the culture medium was almost similar to that observed after a serum-shock: *cry1* mRNA level peaked at Zeitgeber time 8 (ZT8) to ZT10 (where ZT0 is time when the light switched on at 0700 h, and ZT12 is time when the light switched off at 1900 h), *rev-erb $\alpha$*  at ZT16–ZT20, and *bmal1* at ZT4 (Fig. 4C). Thus, fenofibrate can act as a Zeitgeber in cell culture and trigger a rhythm of clock gene expression.

To determine whether fenofibrate can also reset the liver clock *in vivo*, the response of *bmal1* and *rev-erb $\alpha$*  genes to fenofibrate treatment was compared in liver of WT and PPAR $\alpha$ <sup>-/-</sup> mice (Fig. 4D). In accordance with our previous results, fenofibrate was also able to markedly induce *bmal1* mRNA levels in control mice but not in PPAR $\alpha$ -deficient mice, suggesting that *bmal1* induction by fenofibrate is effectively mediated by PPAR $\alpha$ . Similarly, as it was previously reported in rat liver and human hepatocytes (30, 43), administration of fenofibrate in WT mice significantly increased


the *rev-erb $\alpha$*  mRNA levels. As expected, *rev-erb $\alpha$*  expression in PPAR $\alpha$  knockout mice was not induced by fenofibrate, confirming that murine *rev-erb $\alpha$*  induction by fenofibrate is mediated by PPAR $\alpha$ . Taken together, these data strongly suggest that fenofibrate can alter the endogenous rhythm of the liver peripheral clock and *in vivo*.

### PPAR $\alpha$ Is a Direct Regulator of *bmal1* Expression


To specify the regulation of *bmal1* gene expression by PPAR $\alpha$  suggested by our analysis of *bmal1* expression in the PPAR $\alpha$ <sup>-/-</sup> mice, we analyzed the interaction of PPAR $\alpha$  with the *bmal1* promoter. First, we performed a bioinformatic research of potential PPAR $\alpha$  binding sites (PPREs) on the *bmal1* promoter region using the Nubiscan software. Two major PPREs were predicted at the positions –1519 (+, TGGACATGGGTCA) and –4943 (–, AGGGCTGAGGACA), the start site corresponding to the one identified in mouse testis (52). To evaluate whether PPAR $\alpha$  binds to the *bmal1* gene promoter *in vivo*, the occupancy of the potential PPRE binding sites by PPAR $\alpha$  was analyzed using chromatin immunoprecipitation (ChIP) assays performed on hepatocyte DNA using an anti-PPAR $\alpha$  antibody (Fig. 5A). As previously shown *in vitro*, the DNA encompassing the *rev-erb $\alpha$*  Rev-DR2 site (position –45) was precipitated *in vivo* by the anti-PPAR $\alpha$  antibody after fibrate treatment (Fig. 5A, lower panel, lanes 7 and 9), in accordance with a PPAR $\alpha$ -Rev-erb $\alpha$  cross talk through competition for binding to the same Rev-DR2 site (43). Moreover, an amplification product was observed when the same DNA samples were PCR-amplified using primers covering the PPRE located at the position –1519 in the *bmal1* promoter (Fig. 5A, upper panel, lanes 7 and 9). No amplification product was obtained using primers flanking the site at the position –4943 (middle panel). These data further demonstrate that in mice *in vivo*, PPAR $\alpha$  directly binds to the PPRE site located at the position –1519 of the *bmal1* promoter in peripheral oscillators and therefore that *bmal1* is a direct PPAR $\alpha$  target gene.

### PPAR $\alpha$ mRNA Expression Is Severely Down-Regulated in the Liver of *bmal1*<sup>-/-</sup> Mice

To test the hypothesis that the regulation of circadian PPAR $\alpha$  expression involves *bmal1* in the liver, we analyzed the daily accumulation of PPAR $\alpha$  mRNA in the liver of *bmal1*<sup>-/-</sup> mice by quantitative PCR. PPAR $\alpha$  mRNA expression is rhythmic in *bmal1*<sup>-/-</sup> mice with a maximum around ZT12 as observed in WT mice, whereas the amplitude of the peak of expression is significantly dampened (Fig. 5B). This result suggests that BMAL1 is an upstream regulator of PPAR $\alpha$  gene and is consistent with the recent observation that PPAR $\alpha$  expression is also CLOCK dependent (27).



**Fig. 4.** Effect of Serum Shock or Fenofibrate on Clock Gene Expression  
 A, Accumulation of *PPAR $\alpha$*  ( $\blacklozenge$ ), *cry1* ( $\blacktriangle$ ), and *rev-erb $\alpha$*  ( $\blacksquare$ ) mRNAs in Rat-1 fibroblasts shocked with 50% of horse serum. *PPAR $\alpha$*  accumulation in absence of horse serum ( $\blacksquare$ ) is reported as a control of noninduction. The presented values are means  $\pm$  SEM of duplicates of the same reaction for three different experimental points. B, Accumulation of *PPAR $\alpha$*  mRNAs in Rat-1 fibroblasts shocked with 50  $\mu$ M fenofibrate ( $\blacksquare$ ) or with vehicle ( $\blacklozenge$ ) as controls. The presented values are means  $\pm$  SEM of duplicates of the same reaction for three independent experimental points. C, Accumulation of *bmal1*, *cry1*, and *rev-erb $\alpha$*  mRNAs in Rat-1 fibroblasts shocked with fenofibrate ( $\blacksquare$ ) or with vehicle ( $\blacklozenge$ ) as controls. The presented values are means  $\pm$  SEM of duplicates of three independent experimental points. D, Circadian accumulation of *bmal1* and *rev-erb $\alpha$*  mRNAs in liver of WT and *PPAR $\alpha$* <sup>-/-</sup> mice treated ( $\blacksquare$ ) or not ( $\square$ ) with fenofibrate. In this experiment, mice were treated for 2 wk with fenofibric acid (vehicle DMSO) mixed in the drinking water at the final concentration of 7 mM. Control animals were treated with the vehicle in the drinking water. Real-time PCR was used to determine transcript levels. Transcript level values are displayed as relative quantity (RQ) after normalization to the noncyclic 36B4 expression levels in the same sample. Results are expressed as the means  $\pm$  SEM of values from eight animals for both genotypes at each time point.



**Fig. 5.** Reciprocal Regulation of BMAL1 and PPAR $\alpha$


A, ChIP of the potential *bmal1* and *rev-erb $\alpha$*  PPRE elements with PPAR $\alpha$  antibodies. WT and PPAR $\alpha$ <sup>-/-</sup> mice (n = 3 for both genotypes) were fed for 5 d with either Wy14,643 or vehicle (V). ChIP of liver extracts was performed with a PPAR $\alpha$  antibody (PPAR-Ab) and analyzed by PCR for enrichment of the PPRE element of the *bmal1* promoter (*top panels*) and of the *rev-erb $\alpha$*  promoter Rev-DR2 (*bottom panel*). p.i., Preimmune serum; Input, nonprecipitated genomic DNA. B, Daily expression of PPAR $\alpha$  mRNAs in WT (◆) and *bmal1* mutant mice (■) using real-time PCR. Transcript levels were normalized against the noncyclic 36B4 transcript level in the same sample. Results are expressed as the means  $\pm$  SEM of values from three to five animals at each time point. Statistically significant differences between WT and deficient mice are indicated by an asterisk ( $P < 0.0001$ ).

## DISCUSSION

### A New Regulatory Feedback Loop Involved BMAL1 and PPAR $\alpha$ in Peripheral Clocks

Molecular dissection of the mechanisms by which the clock oscillating system is controlled remains one of the most important challenges to assess the importance of the circadian regulation in diverse physiological and metabolic processes in mammals. In the present study, we have integrated the PPAR $\alpha$  gene and its protein into a new positive-regulatory feedback loop in the liver. A model summarizing our main results is shown in Fig. 6. We show that PPAR $\alpha$  plays an important role in the endogenous rhythmic property of peripheral clocks *in vivo*, whereas in the central clock PPAR $\alpha$  deficiency does not alter the circadian expression of clock genes. These clock gene expressions are only affected in the liver of PPAR $\alpha$ -deficient mice. First, PPAR $\alpha$  is not essential to drive the central cir-

cadian system because PPAR $\alpha$ <sup>-/-</sup> mice entrained normally without arrhythmic behavior in constant darkness and displayed no alteration in the amplitude and the phase of circadian expression of the clock genes (*bmal1*, *per2*, *per3*, *cry2*, and *rev-erb $\alpha$* ) compared with WT mice. The lack of PPAR $\alpha$ -dependent clock regulation *in vivo* in the SCN can be due either to the absence of a circadian PPAR $\alpha$  function in the central clock or to the presence of another isotype of PPAR ( $\beta$ ,  $\gamma$ ) or other nuclear receptors that are able to bind to the PPRE site and to exert a compensatory effect. Second, the expression of *bmal1*, an essential gene of the molecular oscillator, is drastically reduced in the liver of PPAR $\alpha$ -deficient mice. This regulation of *bmal1* transcription by PPAR $\alpha$  is likely through a direct binding of PPAR $\alpha$  to the PPRE element located at the position -1519 in the *bmal1* promoter. In addition to the established transactivation of the circadian promoter *rev-erb $\alpha$*  by PPAR $\alpha$  (43), our data largely support the view that PPAR $\alpha$  also associates with the



**Fig. 6.** Model of Cross Talk between PPAR $\alpha$  and Circadian Pathways Depicting the Control of Circadian Regulation by PPAR $\alpha$  in Peripheral Clocks

In mammals, circadian rhythms are generated by the main pacemaker located in the SCN of the hypothalamus, which synchronizes the peripheral oscillators and ensure an ordered response of the organism in terms of physiology, metabolism, and behavior to environmental changes. These peripheral clocks can be reset by alternative routes such as feeding time. Herein, we show that PPAR $\alpha$  is entirely reset by feeding time. We propose that *bmal1* transcription is directly positively regulated by PPAR $\alpha$  and that BMAL1 imposes circadian regulation on PPAR $\alpha$  transcription (*bold arrows*). The amplified accumulation of PPAR $\alpha$  under fibrate treatment leads in turn to a higher level of expression of *bmal1* gene (*dashed arrows*).

circadian *bmal1* promoter *in vivo*. Interestingly, PPAR $\alpha$  expression is strongly decreased in the liver of *bmal1*-deficient mice compared with WT mice. Because no significant change in hepatic CLOCK level is found in *bmal1*-deficient mice compared with WT mice (53), this suggests that BMAL1 itself is in return involved in the circadian transactivation of PPAR $\alpha$  gene at the level of peripheral oscillators in mice. Other observations have also revealed that CLOCK is involved in the circadian transactivation of PPAR $\alpha$  (27) and interacts with its partner RXR (34). We propose that *bmal1* is directly and positively regulated by PPAR $\alpha$  and that BMAL1 imposes in return a circadian regulation on PPAR $\alpha$  transcription.

#### PPAR $\alpha$ Expression Is Reset by Feeding in Peripheral Clocks

Liver is known to be the organ reacting most rapidly to the temporal feeding regimen because it plays a dominant role in the metabolism and processing of food components, such as proteins, lipids, and carbohy-

drates. It was clearly established that the circadian gene expression in peripheral hepatic cells is intimately connected to feeding (10, 11) and that PPAR $\alpha$  is involved in food processing and energy homeostasis (45). A connection between circadian gene regulation, metabolism and energy homeostasis was also established. For example, glucose metabolism (e.g. 6-phosphofructokinase-2, aldolase, and glucose phosphate isomerase) is under the control of the circadian time-keeping system (23, 24). Rhythmic expression of numerous enzymes and transcription factors involved in protein and amino acid metabolism (e.g. serine dehydratase, DBP, and 3-hydroxy-3-methylglutaryl coenzyme A reductase) (7, 54), in fat metabolism (e.g. cholesterol 7 $\alpha$ -hydroxylase, PPAR $\alpha$ , HMG-CoA lyase and reductase) (7, 35, 55), or in detoxification process [e.g. steroid 15 $\alpha$ -hydroxylase (Cyp2a4) and coumarin 7-hydroxylase (Cyp2a5), Cyp2e1, Cyp17, and glutathione-S-transferase  $\theta$ 2] (23, 56) was also observed in liver. Herein, we show that the circadian expression of PPAR $\alpha$  is reset by feeding time and that, in the absence of PPAR $\alpha$ , clock

gene expression in liver is entrained by reversed feeding time, suggesting that PPAR $\alpha$  does not play a major role in food resetting. We underline that, after an extended duration of daytime feeding, the food-imposed reversed phase of circadian gene expression in peripheral liver is similar between WT and PPAR $\alpha$ -deficient mice. This is probably due to PPAR $\alpha$ -independent signaling pathway that plays a major role in the phase resetting of circadian gene expression by feeding time.

### Fibrates as a Resetting Signal in Cell Cultures and Peripheral Clocks

Previous studies have shown that PPAR $\alpha$  expression is positively controlled by glucocorticoids and fibrates and negatively by insulin (37–39). Fibrates are also known to up-regulate the expression of *rev-erb $\alpha$*  in the rat liver and in both rat and human primary hepatocyte cultures (30, 43). *In vitro*, PPAR $\alpha$  mRNA was induced in rat fibroblasts culture after both a serum shock and, more interestingly, a fenofibrate treatment. Moreover, fenofibrate efficiently stimulates the rhythmic expression of several clock genes such as *cry1*, *bmal1*, and *rev-erb $\alpha$* . This suggests that fenofibrate is able to entrain rhythmic PPAR $\alpha$  and clock gene expressions in Rat-1 fibroblasts. *In vivo*, *rev-erb $\alpha$*  mRNA level is up-regulated by fenofibrate in the mouse liver of WT animals. Because this up-regulation of *rev-erb $\alpha$*  gene by fibrates is not observed in the PPAR $\alpha$ -deficient mice, we confirm *in vivo* that the induction of *rev-erb $\alpha$*  gene expression by fibrates is mediated by PPAR $\alpha$  at the transcriptional level. Interestingly, the increased accumulation of PPAR $\alpha$  after fibrate treatment leads in turn to a higher level of expression of *bmal1* in WT mice. Other convincing evidence that the fibrate effect on the *bmal1* expression is exerted at the transcriptional level via the PPAR $\alpha$  protein is that fibrates have no effect on the *bmal1* expression in the PPAR $\alpha$ <sup>-/-</sup> mice. Taken together, these results suggest that PPAR $\alpha$  could play a role in integrating chemical signals inside the liver.

### How Circadian Rhythm Might Influence a Fibrate Therapy?

To date, fibrates are clinically used as hypolipidemic drugs that lower plasma cholesterol and triglycerides. They exert their effect by regulating the expression of several key genes implicated in lipid metabolism via PPAR $\alpha$  activation. Interestingly, fibrate therapy represents a cost-effective approach in the clinical management and the prevention of cardiovascular diseases in a growing population suffering from lifestyle-induced metabolic dysfunctions such as obesity, insulin resistance, and diabetes (57). Although important differences in lipid metabolism exist between mice and human, including the function of PPAR $\alpha$  (58, 59), our *in vitro* and *in vivo* studies suggest that the regulation by fibrates of the circadian expression of

clock genes may influence the success of a treatment because it would suggest a potential induction and subsequently a dysfunction of their expression after fibrate supply in patients. The validity of this hypothesis remains to be demonstrated in humans. Therefore, one additional question to address is how fibrate administration at a selected time of the day can impact the efficacy and the success of the treatment. It will be of interest to decipher the molecular mechanisms involved in the circadian expression of clock genes and PPAR $\alpha$ -regulated genes in presence or absence of fibrates to provide new insight in the downstream circadian physiological and cellular processes governed by PPAR $\alpha$  itself. Similarly, such understanding should lead to new strategies for pharmacological manipulation of the human clock to improve the treatment of dyslipidaemias.

In summary, our data indicate that PPAR $\alpha$  is a specific element of the liver oscillatory clock in mammals and plays an important role in integrating signals into the clock machinery. We clearly demonstrate that, *in vivo*, PPAR $\alpha$  is required to maintain normal circadian oscillation of the master clock gene *bmal1* in liver. This regulation occurs via a direct binding of PPAR $\alpha$  on a PPRE located in the *bmal1* promoter. In addition, BMAL1 is an upstream regulator of the PPAR $\alpha$  expression. This finding provides a new regulatory pathway for the circadian system and suggests that some transcription factors may have a specific role in the peripheral clocks. Further studies are now required to determine the exact impact of circadian rhythms on the metabolic processes governed by PPAR $\alpha$  in peripheral organs. This can be addressed by investigating the circadian regulation of PPAR $\alpha$  target genes and some of the downstream targets in the peripheral clock in WT, PPAR $\alpha$ <sup>-/-</sup>, and other clock mutant mice. The complete elucidation of the signaling elements involved in the interactions between central and peripheral clocks and the mechanisms that govern the interplay between metabolism and circadian oscillators will also have a major impact on the circadian field in the next future.

## MATERIALS AND METHODS

### Cell Culture, Serum, and Fibrate Shock

Rat-1 fibroblasts were grown in DMEM supplemented with 5% fetal calf serum and a mixture of penicillin/streptomycin/glutamine (PSG). The serum shock was done as described elsewhere using 50% horse serum (9). For the fibrate shock, the medium was exchanged with DMEM-PSG supplemented with 50  $\mu$ M fenofibrate (Sigma, St. Louis, MO) after the cells reach confluence (time 0). This medium was replaced with fenofibrate-free DMEM-PSG medium after 2 h. At 0, 4, 8, 12, 16, 20, 24 h after shock, cells were lysed and kept at -70 C until RNA extraction. Whole-cell RNAs were extracted using GenElute Mammalian Total RNA Extraction kit from Sigma.

## Animal Experiments

Purebred WT and homozygous *PPAR $\alpha$ <sup>-/-</sup>* mice on an SV129 background were used. All experiments were done with male mice between 6 and 8 wk of age. Animals were kept under a 12L:12D cycle, and food and drinking water were available *ad libitum*, except when indicated otherwise. The experimental protocols of the current research were approved by the rules and regulations of French veterinary services.

## Locomotor Activity Recording

Adult male mice ( $n = 8$  for both WT and *PPAR $\alpha$ <sup>-/-</sup>*) were exposed to 12L:12D cycle for at least 2 wk. For monitoring locomotor activity, mice were housed individually in cages equipped with infrared motion captors placed over the cages and a computerized data acquisition system (Circadian Activity Monitoring System, Institut National de la Santé et de la Recherche Médicale, France). Activity records were analyzed with the Clocklab software package (Actimetrics, Evanston, IL). For each animal, the total duration of activity was determined every 2 or 12 h during the LD cycle and then averaged for WT and knockout mice. Animals were then allowed to free run in constant darkness (DD) for at least 15 d. The endogenous period in DD was subsequently determined using the Clocklab software.

## Circadian Expression of Clock Genes in Wild-Type vs. PPAR $\alpha$ <sup>-/-</sup> Mice

Twelve WT and 12 *PPAR $\alpha$ <sup>-/-</sup>* mice were maintained for a period of 2 wk in a 12L:12D cycle and transferred in DD the day before the kill. Livers and SCN were removed at CT1, CT8, CT14, and CT21; stored at  $-70$  C until RNA extraction; and analyzed by quantitative RT-PCR. The experiment was done twice.

## Restricted Feeding

Mice ( $n = 24$  both for WT and *PPAR $\alpha$ <sup>-/-</sup>*) fed during the day received food when light was on (0700 to 1900 h), whereas mice ( $n = 24$  both for WT and *PPAR $\alpha$ <sup>-/-</sup>*) fed during the night received food from 1900 to 0700 h for 2 wk. Water was freely available over the experimental period. As controls, WT and *PPAR $\alpha$ <sup>-/-</sup>* mice ( $n = 24$  for both) were fed *ad libitum*. Mice were transferred in DD the day before killing. Livers were dissected at indicated circadian times (CT1, CT8, CT14, CT21), stored at  $-70$  C until RNA extraction, and analyzed by quantitative-RT-PCR.

## Fenofibrate Response

In this experiment, mice ( $n = 32$  both for WT and *PPAR $\alpha$ <sup>-/-</sup>*) were treated for 2 wk with fenofibric acid (vehicle DMSO) mixed in the drinking water at the final concentration of 7 mM. Control animals ( $n = 32$  both for WT and *PPAR $\alpha$ <sup>-/-</sup>*) were treated with vehicle in the drinking water. Livers were removed at CT1, CT8, CT14, and CT21, stored at  $-70$  C until RNA extraction, and analyzed by quantitative RT-PCR.

## Analysis of *bmal1*<sup>-/-</sup> Mice

WT and homozygous *bmal1*<sup>-/-</sup> mice on B6 background (both females and males aged from 8–14 wk provided by C. Bradfield, McArdle Laboratory for Cancer Research, University of Wisconsin, Madison, WI) were kept under 12L:12D cycles and fed *ad libitum*. Livers were removed at indicated Zeitgeber time. Reverse-transcribed total RNAs from three to five animals per time point were analyzed by quantitative RT-PCR as described below.

## RNA Extraction and Quantitative RT-PCR

Total RNAs were prepared from cells or from organs according to the manufacturer's instructions (Sigma) and reverse transcribed using random primers and MMLV Reverse Transcriptase (Invitrogen, San Diego, CA). cDNA were then used as template for a quantitative real-time PCR assay using the QuantiTect SYBR Green PCR reagents (Qiagen, Valencia, CA) and the DNA Engine Opticon system (MJ Research, Cambridge, MA). Each couple of oligonucleotides used were designed to hybridize on different exons. The sequences of forward and reverse primers were as follows: *bmal1*, forward, 5'-CCAAGAAAGTATGGACACAGACAAA-3'; *bmal1*, reverse, 5'-GCATTCTTGATCCTTCCTTGGT-3'; *cry1*, forward, 5'-CTGGCGTGGGAAGTCATCGT-3'; *cry1*, reverse, 5'-CTGTCCGC-CATTGAGTTCTATG-3'; *cry2*, forward, 5'-TGTCCCTCCTGTGTGGAAGA-3'; *cry2*, reverse, 5'-GCTCCCAGCTTGGCTTGAA-3'; *per1*, forward, 5'-GGAGACCACTGAGAGCAGCAA-3'; *per1*, reverse, 5'-CGCACTCAGGAGGCTGTAGGC-3'; *per2*, forward, 5'-ATGCTCGCCATCCACAAGA-3'; *per2*, reverse, 5'-GCGGAATCGAATGGGAGAAT-3'; *per3*, forward, 5'-GGCGTTCTACGCGCACACTGC-3'; *per3*, reverse, 5'-CGCTGGTGCACATTCATACTGCG-3'; *ppar $\alpha$* , forward, 5'-CGCTATGAAGTTCAATGCCTT-3'; *ppar $\alpha$* , reverse, 5'-TGC-AACTTCTCAATGTAGCC-3'; *rev-erb $\alpha$* , forward, 5'-CATGG-TGCTACTGTGTAAGGTGTGT-3'; *rev-erb $\alpha$* , reverse, 5'-CAC-AGGCGTGCCTCCATAG-3'; *36B4*, forward, 5'-ACCTCCT-TCTCCAGGCTTT-3'; and *36B4*, reverse, 5'-CCCACCTTG-TCTCCAGTCTTT-3'.

The efficiency (>95%) and the specificity of the amplification were controlled by generating standard curves and carrying out melting curves and agarose gels of the amplicons, respectively. The relative levels of each RNA were calculated by  $2^{-CT}$  (CT standing as the cycle number in which SYBR Green fluorescence exceeds a constant threshold value) and normalized to the corresponding noncyclic *36B4* RNA levels. The presented values are means  $\pm$  SEM of duplicates of the same reaction for at least three different mice or three experimental points.

The significance of differences was assessed by distribution-free two-way ANOVA. Paired Student's tests were used to compare WT and *PPAR $\alpha$ <sup>-/-</sup>* or *bmal1*<sup>-/-</sup> mice. Results are presented as mean  $\pm$  SEM. Differences were considered significant when  $P < 0.05$ .

## ChIP Assays

ChIP experiments were performed as already described in Ijpenberg et al. (60). Briefly, WT and *PPAR $\alpha$ <sup>-/-</sup>* mice ( $n = 3$ ) were fed for 5 d with either Wy14,643 (50 mg/kg-d) or vehicle. Immunoprecipitation of liver extracts was done using a PPAR $\alpha$  antibody and the immunoprecipitated DNA was PCR amplified using primers flanking either the *rev-erb $\alpha$*  Rev-DR2 (GTGTCACTGGGGC) or potential PPRE (usually AGGT-CANAGGTCA) on the *bmal1* promoter predicted using the computer program NUBISCAN available at the following website: www.nubiscan.unibas.ch (61). An equal volume of nonprecipitated genomic DNA (Input) was amplified as positive control. One fifth of PCR products were separated on an ethidium bromide-stained 2% agarose gel.

## Acknowledgments

We thank J. Samarut and M. Janier for their financial support, and B. Desvergne and G. Triqueneaux for their helpful discussions. We are also grateful to F. J. Gonzalez for the *PPAR $\alpha$ <sup>-/-</sup>* mice.

Received January 30, 2006. Accepted March 15, 2006.

Address all correspondence and requests for reprints to: Vincent Laudet, Centre National de la Recherche Scientifique

Unité Mixte de Recherche 5161, Institut Fédératif de Recherche 128 BioSciences Lyon-Gerland, Laboratoire de Biologie Moléculaire de la Cellule, Ecole Normale Supérieure, 46 allée d'Italie, 69364 Lyon cedex 07, France. E-mail: vincent.laudet@ens-lyon.fr.

This work was supported by Centre National de la Recherche Scientifique; L'École Normale Supérieure de Lyon; Région Rhône-Alpes; Fondation Rhône Alpes Futur; Ministère de l'Éducation Nationale, de la Recherche et de la Technologie; and FP5-OldClock QLK6-CT-2002-02258.

Present address for L.C.: ANIMAGE, Centre National de la Recherche Scientifique Unité Mixte de Recherche 5515, 59 Boulevard Pinel, 69677 Bron cedex, France.

L.C., J.R., O.D.-B., B.R., N.S.T., L.M., F.D., W.W., and V.L. have nothing to declare.

## REFERENCES

- Reppert SM, Weaver DR 2002 Coordination of circadian timing in mammals. *Nature* 418:935–941
- Challet E, Caldeas I, Graff C, Pevet P 2003 Synchronization of the molecular clockwork by light- and food-related cues in mammals. *Biol Chem* 384:711–719
- Albrecht U, Eichele G 2003 The mammalian circadian clock. *Curr Opin Genet Dev* 13:271–277
- Morse D, Sassone-Corsi P 2002 Time after time: inputs to and outputs from the mammalian circadian oscillators. *Trends Neurosci* 25:632–637
- Dunlap JC 1999 Molecular bases for circadian clocks. *Cell* 96:271–290
- Sassone-Corsi P 1998 Molecular clocks: mastering time by gene regulation. *Nature* 392:871–874
- Panda S, Antoch MP, Miller BH, Su AI, Schook AB, Straume M, Schultz PG, Kay SA, Takahashi JS, Hogenesch JB 2002 Coordinated transcription of key pathways in the mouse by the circadian clock. *Cell* 109:307–320
- Kita Y, Shiozawa M, Jin W, Majewski RR, Besharse JC, Greene AS, Jacob HJ 2002 Implications of circadian gene expression in kidney, liver and the effects of fasting on pharmacogenomic studies. *Pharmacogenetics* 12:55–65
- Balsalobre A, Damiola F, Schibler U 1998 A serum shock induces circadian gene expression in mammalian tissue culture cells. *Cell* 93:929–937
- Stokkan KA, Yamazaki S, Tei H, Sakaki Y, Menaker M 2001 Entrainment of the circadian clock in the liver by feeding. *Science* 291:490–493
- Damiola F, Le Minh N, Preitner N, Kornmann B, Fleury-Olela F, Schibler U 2000 Restricted feeding uncouples circadian oscillators in peripheral tissues from the central pacemaker in the suprachiasmatic nucleus. *Genes Dev* 14:2950–2961
- Guo H, Brewer JM, Champhekar A, Harris RB, Bittman EL 2005 Differential control of peripheral circadian rhythms by suprachiasmatic-dependent neural signals. *Proc Natl Acad Sci USA* 102:3111–3116
- Balsalobre A, Marcacci L, Schibler U 2000 Multiple signaling pathways elicit circadian gene expression in cultured Rat-1 fibroblasts. *Curr Biol* 10:1291–1294
- Brown SA, Schibler U 1999 The ins and outs of circadian timekeeping. *Curr Opin Genet Dev* 9:588–594
- Hastings MH, Reddy AB, Maywood ES 2003 A clockwork web: circadian timing in brain and periphery, in health and disease. *Nat Rev Neurosci* 4:649–661
- Reppert SM, Weaver DR 2001 Molecular analysis of mammalian circadian rhythms. *Annu Rev Physiol* 63:647–676
- Lowrey PL, Takahashi JS 2000 Genetics of the mammalian circadian system: photic entrainment, circadian pacemaker mechanisms, and posttranslational regulation. *Annu Rev Genet* 34:533–562
- Ripperger JA, Shearman LP, Reppert SM, Schibler U 2000 CLOCK, an essential pacemaker component, controls expression of the circadian transcription factor DBP. *Genes Dev* 14:679–689
- Ueda HR, Hayashi S, Chen W, Sano M, Machida M, Shigeyoshi Y, Iino M, Hashimoto S 2005 System-level identification of transcriptional circuits underlying mammalian circadian clocks. *Nat Genet* 37:187–192
- Shearman LP, Sriram S, Weaver DR, Maywood ES, Chaves I, Zheng B, Kume K, Lee CC, van der Horst GT, Hastings MH, Reppert SM 2000 Interacting molecular loops in the mammalian circadian clock. *Science* 288:1013–1019
- Marquez S, Crespo P, Carlini V, Garbarino-Pico E, Baler R, Caputto BL, Guido ME 2004 The metabolism of phospholipids oscillates rhythmically in cultures of fibroblasts and is regulated by the clock protein PERIOD 1. *FASEB J* 18:519–521
- Oishi K, Miyazaki K, Kadota K, Kikuno R, Nagase T, Atsumi G, Ohkura N, Azama T, Mesaki M, Yukimasa S, Kobayashi H, Itaka C, Umehara T, Horikoshi M, Kudo T, Shimizu Y, Yano M, Monden M, Machida K, Matsuda J, Horie S, Todo T, Ishida N 2003 Genome-wide expression analysis of mouse liver reveals CLOCK-regulated circadian output genes. *J Biol Chem* 278:41519–41527
- Akhtar RA, Reddy AB, Maywood ES, Clayton JD, King VM, Smith AG, Gant TW, Hastings MH, Kyriacou CP 2002 Circadian cycling of the mouse liver transcriptome, as revealed by cDNA microarray, is driven by the suprachiasmatic nucleus. *Curr Biol* 12:540–550
- Rudic RD, McNamara P, Curtis AM, Boston RC, Panda S, Hogenesch JB, Fitzgerald GA 2004 BMAL1 and CLOCK, two essential components of the circadian clock, are involved in glucose homeostasis. *PLoS Biol* 2:e377
- Turek FW, Joshu C, Kohsaka A, Lin E, Ivanova G, McDearmon E, Laposky A, Losee-Olson S, Easton A, Jensen DR, Eckel RH, Takahashi JS, Bass J 2005 Obesity and metabolic syndrome in circadian Clock mutant mice. *Science* 308:1043–1045
- Inoue I, Shinoda Y, Ikeda M, Hayashi K, Kanazawa K, Nomura M, Matsunaga T, Xu H, Kawai S, Awata T, Komoda T, Katayama S 2005 CLOCK/BMAL1 is involved in lipid metabolism via transactivation of the peroxisome proliferator-activated receptor (PPAR) response element. *J Atheroscler Thromb* 12:169–174
- Oishi K, Shirai H, Ishida N 2005 CLOCK is involved in the circadian transactivation of peroxisome-proliferator-activated receptor  $\alpha$  (PPAR $\alpha$ ) in mice. *Biochem J* 386:575–581
- Shimba S, Ishii N, Ohta Y, Ohno T, Watabe Y, Hayashi M, Wada T, Aoyagi T, Tezuka M 2005 Brain and muscle Arnt-like protein-1 (BMAL1), a component of the molecular clock, regulates adipogenesis. *Proc Natl Acad Sci USA* 102:12071–12076
- Raspe E, Duez H, Gervois P, Fievet C, Fruchart JC, Besnard S, Mariani J, Tedgui A, Staels B 2001 Transcriptional regulation of apolipoprotein C-III gene expression by the orphan nuclear receptor ROR $\alpha$ . *J Biol Chem* 276:2865–2871
- Vu-Dac N, Chopin-Delannoy S, Gervois P, Bonnelye E, Martin G, Fruchart JC, Laudet V, Staels B 1998 The nuclear receptors peroxisome proliferator-activated receptor  $\alpha$  and Rev-erb $\alpha$  mediate the species-specific regulation of apolipoprotein A-I expression by fibrates. *J Biol Chem* 273:25713–25720
- Vu-Dac N, Gervois P, Grotzinger T, De Vos P, Schoonjans K, Fruchart JC, Auwerx J, Mariani J, Tedgui A, Staels B 1997 Transcriptional regulation of apolipoprotein A-I gene expression by the nuclear receptor ROR $\alpha$ . *J Biol Chem* 272:22401–22404
- Hirota T, Okano T, Kokame K, Shirotani-Ikejima H, Miyata T, Fukada Y 2002 Glucose down-regulates Per1

- and Per2 mRNA levels and induces circadian gene expression in cultured Rat-1 fibroblasts. *J Biol Chem* 277:44244–44251
33. Balsalobre A, Brown SA, Marcacci L, Tronche F, Kellendonk C, Reichardt HM, Schutz G, Schibler U 2000 Resetting of circadian time in peripheral tissues by glucocorticoid signaling. *Science* 289:2344–2347
  34. McNamara P, Seo SP, Rudic RD, Sehgal A, Chakravarti D, FitzGerald GA 2001 Regulation of CLOCK and MOP4 by nuclear hormone receptors in the vasculature: a humoral mechanism to reset a peripheral clock. *Cell* 105:877–889
  35. Lemberger T, Saladin R, Vazquez M, Assimakopoulos F, Staels B, Desvergne B, Wahli W, Auwerx J 1996 Expression of the peroxisome proliferator-activated receptor  $\alpha$  gene is stimulated by stress and follows a diurnal rhythm. *J Biol Chem* 271:1764–1769
  36. Ueda HR, Chen W, Adachi A, Wakamatsu H, Hayashi S, Takasugi T, Nagano M, Nakahama K, Suzuki Y, Sugano S, Iino M, Shigeyoshi Y, Hashimoto S 2002 A transcription factor response element for gene expression during circadian night. *Nature* 418:534–539
  37. Lemberger T, Staels B, Saladin R, Desvergne B, Auwerx J, Wahli W 1994 Regulation of the peroxisome proliferator-activated receptor  $\alpha$  gene by glucocorticoids. *J Biol Chem* 269:24527–24530
  38. Steineger HH, Sorensen HN, Tugwood JD, Skrede S, Spydevold O, Gautvik KM 1994 Dexamethasone and insulin demonstrate marked and opposite regulation of the steady-state mRNA level of the peroxisomal proliferator-activated receptor (PPAR) in hepatic cells. Hormonal modulation of fatty-acid-induced transcription. *Eur J Biochem* 225:967–974
  39. Shaw D, Goldman BD 1995 Gender differences in influence of prenatal photoperiods on postnatal pineal melatonin rhythms and serum prolactin and follicle-stimulating hormone in the Siberian hamster (*Phodopus sungorus*). *Endocrinology* 136:4237–4246
  40. Patel DD, Knight BL, Wiggins D, Humphreys SM, Gibbons GF 2001 Disturbances in the normal regulation of SREBP-sensitive genes in PPAR  $\alpha$ -deficient mice. *J Lipid Res* 42:328–337
  41. Gibbons GF, Patel D, Wiggins D, Knight BL 2002 The functional efficiency of lipogenic and cholesterologenic gene expression in normal mice and in mice lacking the peroxisomal proliferator-activated receptor- $\alpha$  (PPAR- $\alpha$ ). *Adv Enzyme Regul* 42:227–247
  42. Kersten S, Seydoux J, Peters JM, Gonzalez FJ, Desvergne B, Wahli W 1999 Peroxisome proliferator-activated receptor alpha mediates the adaptive response to fasting. *J Clin Invest* 103:1489–1498
  43. Gervois P, Chopin-Delannoy S, Fadel A, Dubois G, Kosykh V, Fruchart JC, Najib J, Laudet V, Staels B 1999 Fibrates increase human REV-ERB $\alpha$  expression in liver via a novel peroxisome proliferator-activated receptor response element. *Mol Endocrinol* 13:400–409
  44. Knight BL, Hebbachi A, Hauton D, Brown AM, Wiggins D, Patel DD, Gibbons GF 2005 A role for PPAR $\alpha$  in the control of SREBP activity and lipid synthesis in the liver. *Biochem J* 389:413–421
  45. Desvergne B, Michalik L, Wahli W 2004 Be fit or be sick: peroxisome proliferator-activated receptors are down the road. *Mol Endocrinol* 18:1321–1332
  46. Gachon F, Nagoshi E, Brown SA, Ripperger J, Schibler U 2004 The mammalian circadian timing system: from gene expression to physiology. *Chromosoma* 113:103–112
  47. Canaple L, Kakizawa T, Laudet V 2003 The days and nights of cancer cells. *Cancer Res* 63:7545–7552
  48. Rosbash M 1998 Why the rat-1 fibroblast should replace the SCN as the in vitro model of choice. *Cell* 93:917–919
  49. Yagita K, Tamanini F, van Der Horst GT, Okamura H 2001 Molecular mechanisms of the biological clock in cultured fibroblasts. *Science* 292:278–281
  50. Hastings MH 2005 Circadian biology: fibroblast clocks keep ticking. *Curr Biol* 15:R16–R18
  51. Triqueneaux G, Thenot S, Kakizawa T, Antoch MP, Safi R, Takahashi JS, Delaunay F, Laudet V 2004 The orphan receptor Rev-erb $\alpha$  gene is a target of the circadian clock pacemaker. *J Mol Endocrinol* 33:585–608
  52. Yu Y, Correll PH, Vanden Heuvel JP 2002 Conjugated linoleic acid decreases production of pro-inflammatory products in macrophages: evidence for a PPAR $\gamma$ -dependent mechanism. *Biochim Biophys Acta* 1581:89–99
  53. Kondratov RV, Chernov MV, Kondratova AA, Gorbacheva VY, Gudkov AV, Antoch MP 2003 BMAL1-dependent circadian oscillation of nuclear CLOCK: post-translational events induced by dimerization of transcriptional activators of the mammalian clock system. *Genes Dev* 17:1921–1932
  54. Ogawa H, Ansai Y 1995 Diurnal rhythms of rat liver serine dehydratase, D-site binding protein, and 3-hydroxy-3-methylglutaryl coenzyme A reductase mRNA levels are altered by destruction of the suprachiasmatic nucleus of the hypothalamus. *Arch Biochem Biophys* 321:115–122
  55. Lavery DJ, Schibler U 1993 Circadian transcription of the cholesterol 7 $\alpha$  hydroxylase gene may involve the liver-enriched bZIP protein DBP. *Genes Dev* 7:1871–1884
  56. Lavery DJ, Lopez-Molina L, Margueron R, Fleury-Olela F, Conquet F, Schibler U, Bonfils C 1999 Circadian expression of the steroid 15 $\alpha$ -hydroxylase (Cyp2a4) and coumarin 7-hydroxylase (Cyp2a5) genes in mouse liver is regulated by the PAR leucine zipper transcription factor DBP. *Mol Cell Biol* 19:6488–6499
  57. Vosper H, Khoudoli GA, Graham TL, Palmer CN 2002 Peroxisome proliferator-activated receptor agonists, hyperlipidaemia, and atherosclerosis. *Pharmacol Ther* 95:47–62
  58. Palmer CN, Hsu MH, Griffin KJ, Raucy JL, Johnson EF 1998 Peroxisome proliferator activated receptor- $\alpha$  expression in human liver. *Mol Pharmacol* 53:14–22
  59. Macdonald N, Holden PR, Roberts RA 1999 Addition of peroxisome proliferator-activated receptor  $\alpha$  to guinea pig hepatocytes confers increased responsiveness to peroxisome proliferators. *Cancer Res* 59:4776–4780
  60. Ijpenberg A, Tan NS, Gelman L, Kersten S, Seydoux J, Xu J, Metzger D, Canaple L, Chambon P, Wahli W, Desvergne B 2004 In vivo activation of PPAR target genes by RXR homodimers. *EMBO J* 23:2083–2091
  61. Podvinec M, Kaufmann MR, Handschin C, Meyer UA 2002 NUBIScan, an *in silico* approach for prediction of nuclear receptor response elements. *Mol Endocrinol* 16:1269–1279